Int. J. Environ. Res. Public Health 2014, 11, 9156-9170; doi:10.3390/ijerph110909156

OPEN ACCESS

International Journal of Environmental Research and Public Health ISSN 1660-4601 www.mdpi.com/journal/ijerph

Article

A Dose-Response Relationship between Organic Mercury Exposure from Thimerosal-Containing Vaccines and Neurodevelopmental Disorders

David A. Geier¹, Brian S. Hooker², Janet K. Kern^{1,3}, Paul G. King⁴, Lisa K. Sykes⁴ and Mark R. Geier^{1,*}

* Author to whom correspondence should be addressed; E-Mail: mgeier@comcast.net;

Tel.: +1-301-989-0548; Fax: +1-301-989-1543.

Received: 12 July 2014; in revised form: 7 August 2014 / Accepted: 26 August 2014 / Published: 5 September 2014

Abstract: A hypothesis testing case-control study evaluated concerns about the toxic effects of organic-mercury (Hg) exposure from thimerosal-containing (49.55% Hg by weight) vaccines on the risk of neurodevelopmental disorders (NDs). Automated medical records were examined to identify cases and controls enrolled from their date-of-birth (1991–2000) in the Vaccine Safety Datalink (VSD) project. ND cases were diagnosed with pervasive developmental disorder (PDD), specific developmental delay, tic disorder or hyperkinetic syndrome of childhood. In addition, putative non-thimerosal-related outcomes of febrile seizure, failure to thrive and cerebral degenerations were examined. The cumulative total dose of Hg exposure from thimerosal-containing hepatitis B vaccine (T-HBV) administered within the first six months of life was calculated. On a per microgram of organic-Hg basis, PDD (odds ratio (OR) = 1.054), specific developmental delay (OR = 1.035), tic disorder (OR = 1.034) and hyperkinetic syndrome of childhood (OR = 1.05) cases were significantly more likely than controls to receive increased organic-Hg exposure. By contrast, none of the non-thimerosal related outcomes were significantly more likely than the controls to have received increased organic-Hg exposure. Routine childhood vaccination may be an important public health tool to reduce infectious disease-associated morbidity/mortality, but the present study significantly associates organic-Hg exposure from T-HBV with an increased risk of an ND diagnosis.

Keywords: attention deficit; autism; ethylmercury; merthiolate; thiomersal